



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

rate for the same week of last year, in which it amounted to only 14.8 per thousand. Of the large German towns and cities, more than a third showed less favorable health conditions than Berlin, including Munich, Nuremberg, Carlsruhe, Frankfurt-on-the-Main, Magdeburg, Breslau, Konigsberg, Rixdorf (with 18.6). On the other hand, the following cities showed more favorable health conditions than Berlin, namely: Hamburg, Stuttgart, Cologne, Dusseldorf, Brunswick, Hanover, Charlottenburg (with 13.3), and Schoneberg (with 11.1). The rate of mortality in London was also lower than the Berlin figure, that of Paris and Vienna being higher than the rate of this city. The infant death rate, amounting to 4.1 per year and mille, was higher than the Hamburg rate, but lower than the Munich and Leipzig figure. There were registered 123 deaths from phthisis pulmonalis, 122 deaths from acute diseases of the respiratory organs, 41 deaths from cancer, 16 deaths from scarlet fever, 7 deaths from measles, 3 deaths from diphtheria, and 3 deaths from influenza. Finally, 18 persons died by violence.

#### GUATEMALA.

##### *Report from Livingston, fruit port.*

Acting Assistant Surgeon Peters reports as follows: Week ended June 3, 1905. Present officially estimated population, 3,500; no deaths reported; prevailing diseases malarial; general sanitary condition of this port and the surrounding country during the week, good.

Bills of health were issued to the following-named vessels:

Date.	Vessel.	Number of crew.	Number of passengers from this port.	Number of passengers in transit.	Pieces of baggage disinfected.
May 30	Olympia.....	41	.....	.....	.....
June 1	Belize.....	18	.....	.....	.....

The steamer *Olympia* cleared from Puerto Barrios.

#### HAWAII.

##### *Death from plague at Waipahu.*

HONOLULU, June 21, 1905.

WYMAN, Washington:

There was a death from plague at Waipahu near Honolulu on June 20.

COFER.

#### HONDURAS.

##### *Report from Ceiba, fruit port.*

Acting Assistant Surgeon Robertson reports as follows: Week ended June 10, 1905. Present officially estimated population, about 4,000; one death; prevailing diseases, malarial, mostly mild type, but a few severe cases; general sanitary condition of this port and the surrounding country during the week, good.